CHERE
CENTRE FOR HEALTH ECONOMICS
RESEARCH AND EVALUATION

UNIVERSITY OF
TECHNOLOGY SYDNEY

UTS:CHERE

The Centre for Health Economics Research and Evaluation (CHERE) was established in 1991. CHERE is a centre of excellence in health economics and health services research. It is a joint Centre of the Faculties of Business and Nursing, Midwifery and Health at the University of Technology, Sydney, in collaboration with Central Sydney Area Health Service. It was established as a UTS Centre in February, 2002. The Centre aims to contribute to the development and application of health economics and health services research through research, teaching and policy support. CHERE's research program encompasses both the theory and application of health economics. The main theoretical research theme pursues valuing benefits, including understanding what individuals value from health and health care, how such values should be measured, and exploring the social values attached to these benefits. The applied research focuses on economic and the appraisal of new programs or new ways of delivering and/or funding services. CHERE's teaching includes introducing clinicians, health services managers, public health professionals and others to health economic principles. Training programs aim to develop practical skills in health economics and health services research. Policy support is provided at all levels of the health care system by undertaking commissioned projects, through the provision of formal and informal advice as well as participation in working parties and committees.

University of Technology, Sydney
City campus, Haymarket
PO Box 123 Broadway NSW 2007
Tel: +61 2 9514 4720
Fax: + 61 2 9514 4730
Email: mail@chere.uts.edu.au
www.chere.uts.edu.au

Use of Discrete Choice Experiments in health economics: An update of the literature*

Rochelle Guttmann[1], Ryan Castle[1] & Denzil G. Fiebig[1,2]

CHERE WORKING PAPER 2009/2

1. School of Economics
University of New South Wales

2. Centre for Health Economics Research and Evaluation
Faculty of Business
University of Technology, Sydney

First Version: 17 March 2009
Current Version: 17 March 2009

Abstract

The vast majority of stated preference research in health economics has been conducted in the random utility model paradigm using discrete choice experiments (DCEs). Ryan and Gerard (2003) have reviewed the applications of DCEs in the field of health economics. We have updated this initial work to include studies published between 2001 and 2007. Following the methods of Ryan and Gerard, we assess the later body of work, with respect to the key characteristics of DCEs such as selection of attributes and levels, experimental design, preference measurement, estimation procedure and validity. Comparisons between the periods are undertaken in order to identify any emerging trends.

* This research was supported by the NHMRC through a Program Grant.

Use of Discrete Choice Experiments in health economics: An update of the literature*

Rochelle Guttmann[1], Ryan Castle[1] and Denzil G. Fiebig[1,2]

[1]School of Economics, University of New South Wales
[2]CHERE, University of Technology, Sydney

Version: 17 March, 2009

Abstract

The vast majority of stated preference research in health economics has been conducted in the random utility model paradigm using discrete choice experiments (DCEs). Ryan and Gerard (2003) have reviewed the applications of DCEs in the field of health economics. We have updated this initial work to include studies published between 2001 and 2007. Following the methods of Ryan and Gerard, we assess the later body of work, with respect to the key characteristics of DCEs such as selection of attributes and levels, experimental design, preference measurement, estimation procedure and validity. Comparisons between the periods are undertaken in order to identify any emerging trends.

* This research was supported by the NHMRC through a Program Grant.

Contents

Objective *3*
Method *3*
Results *3*
Background 4
Attribute selection and level setting 6
Experimental design and choice sets 8
Measurement of preferences, response rate and comprehension 10
Estimation and validity 12
Conclusion *13*
References *13*
Appendix: Studies review *14*
January 2001 to June 2004 14
July 2004 to August 2007 16

Objective

The aim of this report is to update a previous review (Ryan and Gerard 2003) of discrete choice experiments (DCEs) applied to a health economics context. Studies published from January 2001 are assessed and evaluated with respect to the key characteristics of a DCE such as selection of attributes and levels, experimental design, preference measurement, estimation procedure and validity. This paper is itself an update of Belkar and Fiebig (2004) and so comparisons are made between three periods.

Method

For full details of the methodology applied, see Ryan and Gerard (2003). In brief, a literature search was conducted to identify published studies using DCEs with a health economics application. The following databases were used: Medline, Embase, Social Science Citation Index, PsychINFO, and EconLIT. Note that the HEALTHSTAR and Health Management Information Consortium databases used in the Ryan and Gerard study were unavailable. This is not believed to present a serious problem because almost all the papers reviewed were found in more than one database. The subsequent terms were used to locate suitable studies: 'conjoint', 'conjoint analysis', 'conjoint measurement', 'conjoint studies', 'conjoint choice experiments', 'part-worth utilities', 'functional measurement', 'paired comparisons', 'pairwise choices', 'discrete choice experiments', 'discrete choice modelling', 'discrete choice conjoint experiments', and 'stated preference'. In addition, papers were identified from the bibliographies of the included studies and selected websites of academic institutions (primarily centres specialising in health economics) to identify discussion papers and conference papers. Only experimental studies grounded in random utility theory (RUT), written in English, applied to health care (as opposed to solely regarding methodology or theory) and based on choice-based response data (as opposed to rating or ranking exercises) were included.

Results

Table 0: History of this review

Stage	*Period Start*	*Period End*	*Months*	*Papers*	*Authors*
I	1990	2000	~120	34	Ryan and Gerard
II	January 2001	June 2004	42	25	Belkar and Fiebig
III	July 2004	August 2007	37	80	Castle and Fiebig

Ryan and Gerard (2003) suggest that stated preference experiments with survey instruments eliciting rankings can be translated into discrete choice experiments for the purposes of

analysis. However they did not find any studies which used this method and our experience of ranking-based studies was the same. All the included studies were based on RUT. If a study's focus was chiefly methodological but an original experiment was conducted as a demonstration, discretion was used in the decision of whether to include it.

January 2001 to June 2004

The literature search generated 55 studies (excluding duplicates) based on a health economics framework, the majority being located through the terms 'conjoint analysis', 'discrete choice experiments' or 'stated preference'. Of these, 50 full references were obtained and through a careful reading 29 of these were excluded. The excluded studies were either solely concerned with methodology or theory of DCEs, used ranking or rating based conjoint analysis, adaptive conjoint analysis, or an earlier publication based on the same experiment was already included in this review or was conducted prior to 2001. The remaining 21 studies were included. A further 7 available studies were identified from bibliographies and academic websites, resulting in a total of 28 studies for analysis (see appendix 1). However, 3 of these articles (marked with an asterisk in the appendix) were included in Ryan and Gerard (2003) and so only 25 studies will be included in the analysis of DCEs produced since 2001.

July 2004 to August 2007

The third period included a far greater number of relevant papers. There were 102 papers which cited the 25 papers included in the previous period. In addition, out of the database search 254 papers were selected from several thousand initial search results leaving a total of 307 unique papers.

For the third period a slightly different approach was taken for the database search. After examining the search databases we discovered that reliable results could be obtained by using broader phrases which enclosed several of the existing phrases. For instance 'discrete choice' encloses 'discrete choice experiment'. Any result found by the latter phrase will also be found by the former. Based on this, we restricted the search phrases to 'conjoint', 'functional measurement', 'paired comparison', 'stated preference' and 'discrete choice'. Since the results from these phrases needed to be exhaustively reviewed by hand, using additional terms did not discover any additional papers. Another issue discovered is that the search databases treat plural and singular forms of the same search term differently (in contrast to the case of web search engines). Using a singular term resulted in a larger number of results, but did not include results where the term was only used in plural form. This should not reduce the proportion of relevant papers found in the third period, but if the selected search terms were followed strictly it is possible that some papers were missed in earlier periods.

From the initial list of 307 papers, a reading of the abstracts eliminated a further 167 papers leaving 140 papers to read thoroughly. Full-text copies of six of these papers could not be located and were excluded. A further 54 papers were excluded based on careful reading, leaving the 80 relevant papers covered here.

Background

Table 1 summarises the background details of the DCE studies in the period of 1990-2000 (reproduced from Ryan and Gerard (2003)), from January 2001 to June 2004 and from July 2004 to August 2007. The number of DCE studies produced has clearly increased

dramatically since the 1990s. The UK remains the largest producer of DCE studies, however the country of origin has diversified markedly. In the past few years a number of studies have been conducted in low income nations, such as China, Vietnam, Ghana and South Africa. Research conducted in these countries faces challenges particular to DCEs including problems measuring willingness-to-pay (WTP) for respondents with dramatically different incomes. The US has also become more involved, maintaining its share of DCE research even as the field expands rapidly.

Table 1: Background of DCEs

Item	*Category*	*Period*					
		1990-2000		2001-2004		2005-2007	
Total		34	100%	25	100%	80	100%
Country	UK	20	59%	10	40%	24	30%
	Australia	6	18%	5	20%	8	10%
	Denmark	0	0%	5	20%	4	5%
	USA	7	21%	3	12%	15	19%
	Other	1	3%	2	8%	29	36%
Year	1990-1997	8	24%				
	1998-1999	13	38%				
	2000	13	38%				
	2001			2	8%		
	2002			10	40%		
	2003			10	40%		
	2004 [a]			3	12%	13	16%
	2005					23	29%
	2006					33	41%
	2007 [b]					11	14%
Area	Economic evaluation	17	50%	22	88%	53	66%
	Insurance plans	5	15%	0	0%	3	4%
	Other	12	35%	3	12%	24	30%
Source [c]	Patient	10	29%	12	48%	41	51%
	Community	11	32%	4	16%	26	33%
	Other	13	38%	11	44%	22	28%

a. Period from 2001 to 2004 includes papers published prior to July 2004
b. Until August 2007
c. Some papers published after 2001 used more than one preference source so columns sum to more than 100%

The vast majority of studies focused on the economic evaluation of health care products. However, other topics of interest included insurance plans and labour issues (e.g. evaluation of job characteristics of health care professionals). Given the considerable proportion of surveys that evaluated preferences for health care products it is no surprise that a large number of studies surveyed patients. In fact, analysing patients' preferences appears to be a growing trend, with less than a third of studies using patients from 1990 to 2000 compared to half of the studies after 2001. Despite the field's growth, research investigating insurance preferences is very uncommon. The three insurance papers we did discover were related to preferences for national health insurance systems and not particular insurance policies.

When evaluating the benefits of a health product the elicitation of doctors' preferences can be as informative as that of a patient, as it is often doctors' opinions that influence patients' decisions. Given this, several studies did use health care professionals such as GPs and nurses. Obtaining preferences from health care professionals does appear to becoming less common however. In the 2004-2007 period only 14 papers (18%) surveyed health professionals. This trend is matched by discussions within the literature emphasising the importance of understanding patient preferences. In fact, one of the claimed benefits of DCEs is that they facilitate shifting decision-making control from professionals to patients. They allow effective aggregation of patient preferences where previously decisions might chiefly have been based upon opinions of experts.

Ryan and Gerard's original paper did not record any papers that surveyed two populations, but this is a fairly common approach. Although, clearly the distinction between two populations and one population differentiated by some attribute may be fine. Approximately 10% of research papers between 2001 and 2007 surveyed two populations. This was typically a comparison between patients and health professionals or occasionally between patient and community preferences. Explicitly surveying two distinct groups is only one method of comparing sub-populations and it was more common to delineate groups using characteristics of the respondent, such as gender or age. While it was rare for researchers to interact experimental attributes they frequently interacted attribute levels with demographic characteristics. This is no doubt partly because interacting attributes required more complex experimental design, but it was also rare to find a theoretical justification for interacting attributes with each other.

Attribute selection and level setting

The majority of studies from 2001 used existing literature in conjunction with focus groups and consultation with experts and/or patients to decide which attributes were important for decision-making. Although some studies used detailed methods to ensure the levels used encompassed a comprehensive range, some appeared to choose a far more arbitrary range.

Table 2 summarises various aspects of attribute selection, namely the number and nature of attributes. The number of attributes used in the studies published between 1990 and 2000 ranged from 2 to 24 whereas the range was much smaller, only 3 to 11, for the period 2001-2004. In both periods the majority of studies contained six or fewer attributes with a mode of six. This tendency continued in the third stage, with over half the studies having from 4 to 6 attributes. Although there appears to be an assumption that a large number of attributes make the task too complicated, only one study in the second period and no studies in the third period assessed the reliability of responses given a change in the number of attributes (from

three to four). In the third period it was common for study designs to follow a specific earlier study or one of several well-known methodological papers (mostly from Louviere, Zwerina or Ryan). Issues such as the number of attributes and the number of levels were rarely discussed in detail and justification for a design was largely deferred to other papers. However, a number of papers did conduct pilot studies and in some cases changed the survey design based on difficulties they encountered.

Table 2: Attribute selection

Item	*Category*	*Period*					
		1990-2000		2001-2004		2005-2007	
Number of attributes [a]	2-3	5	15%	3	12%	18	23%
	4-5	10	29%	11	42%	25	31%
	6	9	26%	6	23%	19	24%
	7-9	4	12%	3	12%	10	13%
	10+	6	18%	3	12%	8	10%
Attributes covered [b]	Monetary measure	19	18%	13	19%	56	12%
	Time	25	24%	13	19%	49	11%
	Risk	12	11%	13	19%	49	11%
	Health status	19	18%	7	10%	101	22%
	Health care	28	26%	7	10%	123	27%
	Other	3	3%	15	22%	73	16%

a. Totals more than 25 for 2001-2004 as one study included two sets of attributes
b. Totals more than the number of studies as each study uses many attributes

In the second period just over half the studies incorporated a monetary attribute such as the cost of a treatment, travel costs or income. The proportion increased slightly in the third period and was included in virtually all cases where a cost attribute was at all realistic. The inclusion of a monetary measure allows for estimation of willingness-to-pay, which is valuable information for new services or for existing services where charging is infeasible. Several papers noted the difficulty in Europe particularly, where patients pay for very little of their own health care. For papers hoping to determine the absolute value of a hypothetical product the experiment required the inclusion of a "opt out" choice. This option was typically the status-quo, which in many cases was a competing form or level of health care provided for free by the government. In these cases determining how valuable a hypothetical product might be is difficult. For instance, one can imagine a DCE where the government hoped to obtain tax-payer valuation of an existing free service, but where "no health care" is not a reasonable "opt-out" choice. This was less of an issue in the US and less developed countries where patients are accustomed to paying for health care.

Where a cost variable was not included, a measure of time frequently served as an indicator of the relative value of other attribute levels. In cases where WTP could be measured this was preferred, but in some contexts it was more appropriate to value trade-offs in one attribute with respect to changes in waiting time. An unusual example of this was found in one paper

which suggested patients were willing to wait an additional two hours to see a GP in exchange for accurate information on the expected wait time. There appears to be a definite trend away from time related attributes. In the first period 24% of all attributes were time-related, but by the third only 11% were. Time-related attributes had also been more common than a cost attribute in the first period, but this preference has reversed. As a measure of the utility obtained from other attributes, cost offers a more natural comparison than time and WTP is used even in contexts where actual user fees would be unlikely. Cost is also more likely to be linearly related to utility than is time, at least in the domains relevant to health care. This assumption is almost always made for both cost and time, but may be more reasonable for cost. Other measures of time were also used, such as travel time and life expectancy but these were rare.

Despite problems communicating the impact of changes in risk levels, a number of papers used an attribute related to risk. Risk was usually measured as the chance of success or failure (e.g. chance of side effects, chance of detection) or as a level of certainty (e.g. certainty of budget expenditure). A number of different techniques were used to describe and explain risks to respondents. Especially for marginal changes in the risk of very unlikely events, respondents struggled to comprehend the distinction. Some studies used graphical representations to characterise risk. Others applied risk probabilities to specific situations that might be familiar to the respondent to give them a more concrete sense of the risk involved. Throughout the literature, health professionals were found to be less risk averse than patients. One paper suggested that this may be mostly due to the fact that health professionals are in a better position to observe what a given risk probability translates to in practice. Perhaps because patients may only get “one roll of the dice”, typical or “mean” welfare outcomes are less meaningful to them.

With respect to the studies produced after 2000, measures of health status include severity of symptoms and illness and quality of life measurements. Many studies incorporated several attributes describing different levels of symptoms and side effects under different treatments. The attributes included in the health care category in Table 2 ranges from the overall health care system available (such as availability of vaccines, financing and choice of hospital), to attributes regarding individual care, such as continuity of care and affability of doctors. When compared to the attributes covered in 1990-2000, there was a large disparity with regard to health status and health care. This, however, could be due to the possibility that the classification of attributes belonging to the health care, health status and consequently the other category, may differ between this paper and that of Ryan and Gerard (2003). Some papers merely aimed to assess the relative importance of a long list of health care or health status attributes. A small number of these papers may have had a large impact on the proportional representation of each attribute type.

Experimental design and choice sets

Table 3 describes various aspects of design choice in DCEs. Of those studies that reported the type of design, virtually all used a fractional factorial design. The large number of possible combinations of attributes rendered it impractical to use a full factorial design for all but four studies in the 1990 to 2000 period. Prior to 2005, over half of the designs examined were generated using software packages, the most popular being SPEED and Orthoplan in SPSS. In the third period, this changed somewhat with growing recognition of the complexity of stated

preference experimental design. Researchers were much more likely to design the experiment themselves, based on progress in the study of optimal research designs. Here, these studies have been recorded as not reporting the source of their design, however the majority of them used some form of design theory to inform the design process. By far the most common reference was to Zwerina et al (1996) who developed a method for constructing D-efficient choice designs.

Table 3: Experimental design and choice sets

Item	*Category*	*Period*					
		1990-2000		2001-2004		2005-2007	
Design Type	Full factorial	4	12%	0	0%	8	10%
	Fractional factorial	25	74%	24	96%	70	88%
	Not reported	5	15%	1	4%	2	3%
Design Source	Software packages	19	66%	14	56%	32	40%
	Catalogue	2	7%	2	8%	4	5%
	Expert	4	14%	5	20%	2	3%
	Not reported	4	14%	4	16%	42	53%
Design Plan	Main effects only	25	74%	13	52%	65	81%
	Main effects and selected two-way interactions	2	6%	6	24%	11	14%
	Full factorial	4	12%	0	0%	3	4%
	Not reported	3	9%	6	24%	1	1%
Allocation of scenarios to choice sets	Binary (yes or no)	2	6%	4	16%	7	9%
	Pairwise (random)	15	44%	13	52%	15	19%
	Pairwise (other)	9	26%	4	16%	32	40%
	Greater than 2 choices	5	15%	4	16%	25	31%
	Not reported	3	9%	0	0%	1	1%

In terms of design plan, over half the studies reviewed in the second period used an orthogonal main effects design although there was a growing tendency to include some two-way interactions. In the third period main effects designs dominated the literature even more, which may be partly due to the greater number of researchers constructing their own designs and looking to Zwerina's preference for orthogonality.

With respect to the construction of choice sets the preferred methods have changed substantially. Pairwise choice sets were used by 68% of studies in the second period and by 80% of the studies in the third period. Increasingly, the process of allocating alternatives to

choice sets has become part of the overall design process. Only 19% of studies randomly paired alternatives in the third period.

Clearly there have been few studies employing more than two alternatives per choice set,. Within this 'greater than two' choice category, choice sets involving a pairwise choice plus an opt-out option (i.e. choose between A, B or neither) are included. In the 2001-2004 period, three studies (12%) incorporated this opt-out structure. In the last period this increased to 17 studies (21%). In some cases the opt-out choice was a hypothetical situation specified by the researcher, but mostly it represented the status quo for the respondent themselves. A number of studies discussed the problems with forcing a choice between two unappealing alternatives and many recognised that an opt-out offered respondents a more realistic experiment. The binary option is akin to the pairwise choice with an opt-out, only with a single alternative presented.

Measurement of preferences, response rate and comprehension

Table 4 suggests a shift towards smaller experiments in 2001-2004 which reversed in 2005-2007. Shorter surveys were thought to be more valid, because respondents were more likely to think about their responses. However, it seems that respondents are willing to answer longer surveys and the understanding they gain during the survey process improves the quality of their responses. Longer surveys also have obvious efficiency benefits. During 2001-2004, a number of studies used two versions of their survey using a different number of choices so as to test whether respondents' decisions changed when presented with more choice sets. None of the studies in the last period did this and none mentioned any problems with survey length. However, only a third of the studies in the final period stated that they had measured how long the survey took and only 16% said they asked respondents about difficulties completing the survey.

The bulk of studies from 1990-2000 used self-complete questionnaires and although computer and internet surveys have increased in popularity, the dominance of traditional methods of survey administration has continued. In the third period, self-complete questionnaires were much less likely to be solicited via mail. This can be seen in the statistics for the number of mail reminders, figures only relevant to postal surveys. It is surprising that internet questionnaires have not become more popular in recent years. It may be attributable to the complexity of constructing DCEs because one of the benefits of the internet is it allows researchers to publish questionnaires online quickly and cheaply. The preference for offering all respondents identical surveys also nullifies a key benefit of the dynamic nature of online surveys. It is also likely that the lack of suitable software plays a role in inhibiting the growth of online DCEs.

Table 4: Measurement of preferences, response rate and comprehension

Item	*Category*	*Period*					
		1990-2000		2001-2004		2005-2007	
Number of choices per respondent	8 or less choices	13	38%	15	60%	33	41%
	9-16 choices	18	53%	7	28%	30	38%
	More than 16	2	6%	3	12%	13	16%
	Not reported	1	3%	0	0%	4	5%
Administration of survey [a]	Self-complete questionnaire	27	79%	17	65%	46	57%
	Interviewer administered	3	9%	8	31%	21	26%
	Computer questionnaire	3	9%	1	4%	7	9%
	Internet questionnaire	-		-		7	9%
	Not Reported	1	3%	0	0%	0	0%
Reminders	None	6	22%	1	6%	6	8%
	1-2 reminders	12	44%	8	47%	13	16%
	More than 2 reminders	0	0%	1	6%	3	4%
	Not Applicable/Not Reported	9	33%	7	41%	58	73%
Response rate (%)	90-100	7	21%	2	8%	14	18%
	70-<90	7	21%	4	16%	18	23%
	50-<70	7	21%	10	40%	11	14%
	30-<50	6	18%	5	20%	11	14%
	<30	2	6%	0	0%	5	6%
	Not reported	5	15%	4	16%	21	26%
Ease of task as perceived by subject	Reported	10	29%	3	12%	13	16%
	Not reported	24	71%	22	88%	67	84%
Completion time	Reported	5	15%	6	24%	26	33%
	Not Reported	29	85%	19	76%	54	68%

a. Totals include two studies where there was an option of an interview or questionnaire

The response rates, although wide-ranging across surveys in all three periods, appear to be fairly constant. They were noticeably lower in the second period, but this hasn't become a trend. In the third period there were many studies that did not sample from a defined population so it was not possible to determine a true response rate. Frequently they reported a response rate of 95% or higher, representing the number of respondents who answered the survey accurately. However, in these cases we marked the response rate as "Not Reported" since the statistic was not meaningful. It is discouraging to see that the number of studies that asked respondents to report on ease of completion has decreased significantly. Moreover,

despite an increase in the number of studies reporting completion time, the figure is still very low.

Estimation and validity

Table 5 summarises the type of econometric model used to analyse the DCE along with a variety of validity measures to test the accuracy and appropriateness of the DCE. Given the extensive use of binary and pairwise choice sets it is no surprise that the probit and logit are the most commonly used model. In particular, the random-effects probit remains a popular choice model, as it allows for a multiple response structure typical of DCE data. In the third period however, some form of a logit model was the most popular, although it was rare for this to be a pure binomial logit model. More frequently, it was a multinomial logit or a customised logit model. One of these in the latter category used a design of only one choice per respondent but the other failed to account for the multiple respondents characterised in their design.

It appears that the definition of "face" validity used by this paper is different from that used by Ryan and Gerard. In the latter two periods the vast majority of studies were categorised as having face validity, but it is a difficult concept to measure. "Theoretical" validity is similarly imprecise and difficult to measure, so comparison across time is probably not meaningful. We have interpreted face validity as conducting pilot studies prior to the actual experiment and modifying surveys where necessary or consulting with experts and respondents in the formulation of the survey. This process was very common (a strong majority conducted pilot studies) in studies published after 2000. This is in contrast to the period prior to 2001 when only two studies (6%) indicated that such tests were undertaken. However, this may well be due to Ryan and Gerard (2003) defining face validity differently, only including those that conducted pre-pilot studies. The bulk of studies in the first and last periods and all the studies in the middle period considered theoretical validity, at least, by verifying expected signs after model estimation.

The majority of studies tested for lexicographic responses. Rationality and willingness to trade are both simple to test for, although the appropriate response to violations of these assumptions isn't clear. In most cases excluding respondents who violated rationality assumptions or did not trade did not substantively affect the results, so studies that tested for these issues typically included these observations anyway. In the first two periods, only one study tested for reliability (i.e. testing more than once over a time period). The proportion of studies testing for reliability increased in the third period. Studies were also more likely to consider convergence of DCE results with other research methods, although this validity criterion is vaguer than testing for reliability over time. There have been some doubts about the general validity of DCEs and this may have prompted greater concern with ability to generalise from DCE results.

Table 5: Estimation and validity

Item	*Category*	*Period*					
		1990-2000		2001-2004		2005-2007	
Econometric model	Probit/ ordered probit	3	9%	1	4%	5	6%
	Random-effects probit	17	50%	17	68%	21	26%
	Logit (incl. RE logit, MNL)	9	26%	7	28%	32	40%
	Other	3	9%	0	0%	19	24%
	Not reported	2	6%	0	0%	3	4%
Validity	Face	2	6%	20	80%	64	80%
	Theoretical	22	65%	25	All	53	66%
	Convergent	4	12%	4	16%	20	25%
	Rationality	18	53%	14	56%	39	49%
	Compensatory Decision Making	12	35%	8	32%	30	38%
	Reliability	1	3%	0	0%	10	13%

Conclusion

The number of discrete choice experiments for application in healthcare is growing rapidly, with 34 studies being analysed between 1990 and 2000 (3 per year), compared with 25 between 2001 and 2004 (7 per year) and 80 between 2004 and 2007 (27 per year).

Despite the large number of healthcare products and issues being evaluated using DCEs, the methods and design employed are quite similar. Most studies used main effects fractional factorial designs generated from software packages for the purpose of economic evaluation. The majority of studies reviewed presented a small number of pairwise choice sets to respondents in the form of a questionnaire. Attributes are usually chosen based on consultations and a literature review and pilot studies are used to refine the surveys. The use of a random-effects model captures the multiple response structure of the experiment and checking for theoretical validity by comparing parameter signs against a priori expectations is de rigeur.

References

Belkar, R. and Fiebig, D.G. (2004), "Use of discrete choice experiments in health economics: An update of the literature", working paper, School of Economics, University of New South Wales.

Ryan M., Gerard K. (2003), "Using discrete choice experiments to value health care programmes: current practice and future research reflections", *Applied Health Economics and Health Policy,* 2 (1), 55-64

Zwerina, Klaus, Joel Huber and Warren Kuhfeld (1996), "A General Method for Constructing Efficient Choice Designs," Available at http://support.sas.com/techsup/technote/ts722e.pdf

Appendix: Studies reviewed

January 2001 to June 2004

1) Aristides, M., Chen, J., Schulz, M., et al. (2002), "Conjoint analysis of a new chemotherapy – Willingness to pay and preference for the features of raltitrexed versus standard therapy in advanced colorectal cancer", *Pharmacoeconomics*, 20 (11), 775-784

2) Bech, M. (2003), "County council politicians' choice of hospital payment scheme: a discrete choice study", *Applied Health Economics and Health Policy*, 2 (4), 225-32

3) Bech, M. (2003), "Politicians' and hospital managers' trade-offs in the choice of reimbursement scheme: a discrete choice experiment", *Health Policy*, 66 (3), 261-275

4) Bishop, A.J., Marteau, T.M., Armstrong, D., et al. (2004), "Women and health care professionals' preferences for Down's syndrome screening tests: a conjoint analysis study", *BJOG- An International Journal of Obstetrics and Gynaecology*, 111, 775-779

5) Bryan, S. and Parry, D. (2002), "Structural reliability of conjoint measurement in health care: an empirical investigation", *Applied Economics*, 34 (5), 561-567

6) Bryan, S., Roberts, T., Heginbotham, C. and McCallum, A. (2002), "QALY-maximisation and public preferences: results from a general population survey", *Health Economics*, 11 (8), 679-693

7) Gerard, K., Shanahan, M. and Louviere, J. (2003), "Using stated preference discrete choice modelling to inform health care decision-making: A pilot study of breast screening participation", *Applied Economics*, 35 (9), 1073-1085

8) Gyrd-Hansen, D. (2003), "Willingness to pay for a QALY", *Health Economics*, 12 (12), 1049-1060

9) Gyrd-Hansen, D. and Slothuus, U. (2002), "The citizen's preferences for financing public health care: a Danish survey", *International Journal of Health Care Finance and Economics*, 2 (1), 25-36

10) Hall, J., King, M., Louviere, J., et al. (2002), "Using stated preference discrete choice modelling to evaluate the introduction of varicella vaccination", *Health Economics*, 11 (5), 457-465

11) Halpern, S.D., Berns, J.S., Israni, A.K. (2004), "Willingness of patients to switch from conventional to daily hemodialysis: Looking before we leap", *American Journal of Medicine*, 116 (9), 606-612

12) Hundley V., Ryan M., Graham W. (2001), "Assessing women's preferences for intrapartum care", *Birth*, 28 (4), 254-263

Hundley, V. and Ryan, M. (2004), "Are women's expectations and preferences for intrapartum care affected by the model of care on offer?", *BJOG- An International Journal of Obstetrics and Gynaecology*, 111 (6), 550-560

13) Lancsar, E.J., Hall, J.P., King, M., et al. (2003), "Using discrete choice experiments to investigate patient preferences for preventative asthma medication", *Paper prepared for the Econometric Society Australasian Meeting*, 2003 9-11 July

14) Longworth, L., Ratcliffe, J. and Boulton, M. (2001), "Investigating women's preferences for intrapartum care: home versus hospital births", *Health and Social Care in the Community*, 9 (6), 404-413

Ratcliffe, J. and Longworth, L. (2002), "Investigating the structural reliability of a discrete choice experiment within health technology assessment", *International Journal of Technology Assessment in Health Care*, 18 (1), 139-144

15) Mark, T.L. and Swait, J. (2003), "Using stated preference modelling to forecast the effect of medication attributes on prescriptions of alcoholism medications", *Value in Health*, 6 (4), 474-482

Mark, T.L. and Swait, J. (2004), "Using stated preference and revealed preference modelling to evaluate prescribing decisions", *Health Economics*, 13 (6), 563-573

16) McKenzie, L., Cairns, J. and Osman, L. (2001), "Symptom-based outcome measures for asthma: the use of discrete choice methods to assess patient preferences", *Health Policy*, 57 (3), 193-204

17) Moayyedi, P., Wardman, M., Toner, J., et al. (2002), "Establishing patient preferences for gastroenterology clinic reorganization using conjoint analysis", *European Journal of Gastroenterology and Hepatology*, 14(4), 429- 433

18) Phillips, K.A., Maddala, T. and Johnson, F.R. (2002), "Measuring preferences for health care interventions using conjoint analysis: An application to HIV testing", *Health Services Research*, 37 (6), 1681-1705

Phillips, K.A., Johnson, F.R. and Maddala, T. (2002), "Measuring what people value: A comparison of "attitude" and "preference" surveys", *Health Services Research*, 37 (6), 1659-1679

Maddala, T., Phillips, K.A. and Johnson, F.R. (2003), "An experiment on simplifying conjoint analysis designs for measuring preferences", *Health Economics*, 12 (12), 1035-1047

19) Ratcliffe, J., Buxton, M., McGarry, T., et al. (2004), "Patients' preferences for characteristics associated with treatments for osteoarthritis", *Rheumatology*, 43 (3), 337-345

20) Ratcliffe, J., Van Haselen, R., Buxton, M., et al. (2002), “Assessing patients’ preferences for characteristics associated with homeopathic and conventional treatment of asthma: a conjoint analysis study”, *Thorax*, 57 (6), 503-508

21)* Ryan, M., Bate, A., Eastmond, C.J. and Ludbrook, A. (2001), “Use of discrete choice experiments to elicit preferences”, *Quality in Health Care*, 10, 155-160 Suppl.

Ryan, M., Bate, A. (2001), “Testing the assumptions of rationality, continuity and symmetry when applying discrete choice experiments in health care”, *Applied Economics Letters*, 8 (1), 59-63

22)* Scott, A. (2001), “Eliciting GPs’ preferences for pecuniary and non-pecuniary job characteristics”, *Journal of Health Economics*, 20 (3), 329-347

23)* Scott, A., Watson, M.S. and Ross, S. (2003), “Eliciting preferences of the community for out of hours care provided by general practitioners: a stated preference discrete choice experiment”, *Social Science and Medicine*, 56 (4), 803-814

Scott, A. (2002), “Identifying and analysing dominant preferences in discrete choice experiments: an application in health care”, *Journal of Economic Psychology*, 23 (3), 383-398

24) Skjoldborg, U.S. and Gyrd-Hansen, D. (2003), “Conjoint analysis. The cost variable: an Achilles’ heel?”, *Health Economics*, 12 (6), 479-491

25) Taylor, S.J. and Armour, C.L. (2003), “Consumer preference for dinoprostone vaginal gel using stated preference discrete choice modelling”, *Pharmacoeconomics*, 21 (10), 721-735

26) Telser, H. and Zweifel, P. (2002), “Measuring willingness-to-pay for risk reduction: an application of conjoint analysis”, *Health Economics*, 11 (2), 129-139

27) Ubach, C., Bate, A., Ryan, M., et al. (2002), “Using discrete choice experiments to evaluate alternative electronic prescribing systems”, *The International Journal of Pharmacy Practice*, 10(3), 191-200

28) Ubach, C., Scott, A., French, F., et al. (2003), “What do hospital consultants value about their jobs? A discrete choice experiment”, *British Medical Journal*, 326 (7404), 1432-1437

July 2004 to August 2007

1) Tsuge T., Kishimoto A. and Takeuchi K. (2005), "A Choice Experiment Approach to the Valuation of Mortality", *Journal of Risk and Uncertainty*, 31 (1), 73-95

2) Lewis, S.M., Cullinane, F.N., Bishop, A.J., Chitty, L.S., Marteau, T.M. and Halliday, J.L. (2005), "A comparison of Australian and UK obstetricians' and midwives' preferences for screening tests for Down syndrome", *Prenatal Diagnosis*, 26, 60-66

3) Lee, A. Gin, T., Lau, A.S.C and Ng, F.F. (2005), "A comparison of patients' and health care professionals' preferences for symptoms during immediate postoperative recovery and the management of postoperative nausea and vomiting", *Anesthesia and Analgesia*, 100 (1), 87-93

4) Jan, M., Fu, T. and Huang, C.L. (2007), "A Conjoint/Logit Analysis of Consumers' Responses to Genetically Modified Tofu in Taiwan", *Journal of Agricultural Economics*, 58 (2), 330-347

5) Peacock, S., Apicella, C., Andrews, L., Tucker, K., Bankier, A., Daly, M.B. and Hopper, J.L. (2006), "A discrete choice experiment of preferences for genetic counselling among Jewish women seeking cancer genetics services", *British Journal of Cancer*, 95, 1448-1453

6) Douglas, H., Normand, C., Higginson, I. and Goodwin, D. (2005), "A New Approach to Eliciting Patients' Preferences for Palliative Day Care: The Choice Experiment Method", *Journal of Pain and Symptom Management*, 29 (5)

7) Telser, H. and Zweifel, P. (2006), "A New Role For Consumers' Preferences In The Provision Of Healthcare", *Economic Affairs*, September 2006

8) Johnson, F.R., Hauber, A.B., Osoba, D., Hsu, M., Coombs, J. and Copley-Merriman, C. (2006), "Are chemotherapy patients' HRQoL importance weights consistent with linear scoring rules? A stated-choice approach", *Quality of Life Research*, 15, 285-298

9) Chattopadhyay, S., Braden, J.B., and Patunru, A. (2005), "Benefits of hazardous waste cleanup: New evidence from survey- and market-based property value approaches", *Contemporary Economic Policy*, 23 (3), 357-375

10) Bishai, D., Brice, R., Girod, I., Saleh, A. and Ehreth, J. (2007), "Conjoint analysis of French and German parents' willingness to pay for meningococcal vaccine", *PharmacoEconomics*, 25 (2), 143-154

11) Skjoldborg, U.S., Lauridsen, J. and Junker, P. (2005), "Conjoint Reliability Measured at the Input and Output Level", *Health Economics Paper*, 2005:7, University of Southern Denmark

12) Kerssens, J.J. and Groenewegen, P.P. (2005), "Consumer preferences in social health insurance", *The European Journal of Health Economics*, 50 (8), 8-15

13) Zweifel, P., Telser, H. and Vaterlaus, S. (2006), "Consumer resistance against regulation: The case of health care", *Journal of Regulatory Economics*, 29, 319-332

14) Mantovani, L.G., Monzini, M.S., Mannucci, P.M., Scalone,L., Villa, M. and Gringeri, A. (2005), "Differences between patients', physicians' and pharmacists' preferences for treatment products in haemophilia: a discrete choice experiment", *Haemophilia*, 11, 589-597

15) Weston, A. and FitzGerald, P. (2004), "Discrete choice experiment to derive willingness to pay for methyl aminolevulinate photodynamic therapy versus simple excision surgery in basal cell carcinoma", *PharmacoEconomics*, 22 (18), 1195-1208

16) Salkeld, G., Solomon, M., Butow, P. and Short, L. (2005), "Discrete-choice experiment to measure patient preferences for the surgical management of colorectal cancer", *British Journal of Surgery*, 92, 742-747

17) Burge, P., Devlin, N., Appleby, J., Rohr, C., and Grant, J. (2004), "Do patients always prefer quicker treatment? A discrete choice analysis of patients' stated preferences in the London Patient Choice Project", *Applied Health Economics and Health Policy*, 3 (4), 183-194

18) Schwappach, D.L.B. and Strasmann, T.J. (2007), "Does location matter? A study of the public's preferences for surgical care provision", *Journal of Evaluation in Clinical Practice*, 13, 259-264

19) Langer, C.J., Fastenau, J.M., Forlenza, J.B., Piech, C.T., et al (2007), "Effectiveness versus convenience: patient preferences for an erythropoietic agent to treat cancer-related anemia", *Current Medical Research and Opinion*, 23 (1), 85-92

20) Watson, V., Ryan, M., Brown, C.T., Barnett, G., Ellis, B.W., and Emberton, M. (2004), "Eliciting preferences for drug treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia", *The Journal of Urology*, 172, 2321-2325

21) Johnson, F.R. and Backhouse, M. (2006), "Eliciting stated preferences for health-technology adoption criteria using paired comparisons and recommendation judgments", *Value in Health*, 9 (5), 303-311

22) Yeung, R.Y.T, Smith, R.D., Ho, L., Johnston, J.M. and Leung, G.M. (2006), "Empirical implications of response acquiescence in discrete-choice contingent valuation", *Health Economics*, 15, 1077-1089

23) Haughney, J. Partridge, M.R., Vogelmeier, C., Larsson, T., Kessler, R., Ståhl, E., Brice, R. and Löfdahl, C. (2005), "Exacerbations of COPD: Quantifying the patient's perspective using discrete choice modelling", *European Respiratory Journal*, 26, 623-629

24) Bech, M., Gyrd-Hansen, D., Kjær, T., Lauridsen, J. and Sørensen, J. (2006), "Graded pairs comparison - Does strength of preference matter? Analysis of preferences for specialised nurse home visits for pain management", *Health Economics*, 16, 513-529

25) Johnson, F.R., Clayton, L.J., Manjunath, R., Hoerger, T.J., Mansfield, C.A. and Zhang, P. (2006), "High-risk individuals' willingness to pay for diabetes risk-reduction programs", *Diabetes Care*, 29 (6), 1351-1356

26) Canh, D.G., Whittington, D., Thoa, L.T.K., Utomo, N., Hoa, N.T., Poulos, C., Thuy, D.T.D., Kim, D., Nyamete, A. and Acosta, C. (2006), "Household demand for typhoid fever vaccines in Hue, Vietnam", *Health and Policy Planning*, 21 (3), 241-255

27) Kjær, T., Gyrd-Hansen, D. and Willaing, I. (2006), "Investigating patients' preferences for cardiac rehabilitation in Denmark", *International Journal of Technology Assessment in Health Care*, 22 (2), 211-218

28) Longo, M.F., Cohen, D.R., Hood, K., Edwards, A., Robling, M., Elwyn, G. and Russell, I.T (2006), "Involving patients in primary care consultations: assessing preferences using discrete choice experiments", *British Journal of General Practice*, 56, 35-42

29) Cunningham, C.E., Deal, K., Neville, A., Rimas, H. and Lohfeld, L. (2006), "Modeling the problem-based learning preferences of McMaster University undergraduate medical students using a discrete choice conjoint experiment", *Advances in Health Sciences Education*, 11, 245-266

30) Yannis, G., Kanellopoulou, A., Aggeloussi, K. and Tsamboulas, D. (2005), "Modelling driver choices towards accident risk reduction", *Safety Science*, 43, 173-186

31) Hojman, P., de Dios Ortuzar, J. and Rizzi, L.I. (2005), "On the joint valuation of averting fatal and severe injuries in highway accidents", *Journal of Safety Research*, 36, 377-386

32) Kjær, T., Bech, M., Gyrd-Hansen, D. and Hart-Hansen, K. (2006), "Ordering effect and price sensitivity in discrete choice experiments: need we worry?", *Health Economics*, 15: 1217-1228

33) Dickie, M. and Messman, V.L. (2006), "Parental altruism and the value of avoiding acute illness: are kids worth more than parents?", *Journal of Environmental Economics and Management*, 48, 1146-1174

34) Aristides, M., Weston, A.R., Fitzgerald, P., Le Reun, C. and Maniadakis, N. (2004), "Patient preference and willingness-to-pay for Humalog Mix25 relative to Humulin 30/70: a multicountry application of a discrete choice experiment", *Value in Health*, 7 (4), 442-454

35) Mahadevia, P.J., Shah, S., Leibman, C., Kleinman, L. and O'Dowd, L. (2004), "Patient preferences for sensory attributes of intranasal corticosteroids and willingness to adhere to prescribed therapy for allergic rhinitis: a conjoint analysis", *Annals of Allergy, Asthma and Immunology*, 93, 345-350

36) Fraenkel, L., Gulanski, B. and Wittink, D. (2006), "Patient treatment preferences for osteoporosis", *Arthritis and Rheumatism*, 55 (5), 729-735

37) Akkazieva, B., Gulacsi, L., Brandtmuller, A., Péntek, M. and Bridges, J.F.P. (2006), "Patients' preferences for healthcare system reforms in Hungary: A conjoint analysis", *Applied Health Economics and Health Policy*, 5 (3), 189-198

38) Alberini, A., Tonin, S., Turvani, M. and Chiabai, A. (2007), "Paying for Permanence: Public Preferences for Contaminated Site Cleanup", *Journal of Risk and Uncertainty*, 34, 155-178

39) Rubin, G., Bate, A., George, A., Shackley, P. and Hall, N. (2006), "Preferences for access to the GP: A discrete choice experiment", *British Journal of General Practice*, 56, 743-748

40) Fraenkel, L., Gulanski, B. and Wittink, D.R. (2006), "Preference for hip protectors among older adults at high risk for osteoporotic fractures", *Journal of Rheumatology*, 33 (10), 2064-2068

41) Coast, J., Salisbury, C., de Berker, D., Noble, A., Horrocks, S., Peters, T.J. and Flynn, T.N. (2006), "Preferences for aspects of a dermatology consultation", *British Journal of Dermatology*, 155, 387-392

42) Wordsworth, S., Skåtun, D., Scott, A. and French, F. (2004), "Preferences for general practice jobs: a survey of principals and sessional GPs", *British Journal of General Practitioners*, 54, 740-746

43) Hanson, K., McPake, B., Nakamba, P. and Archard, L. (2005), "Preferences for hospital quality in Zambia: Results from a discrete choice experiment", *Health Economics*, 14, 687-701

44) Porteous, T., Ryan, M., Bond, C.M. and Hannaford, P. (2006), "Preferences for self-care or professional advice for minor illness: a discrete choice experiment", *British Journal of General Practic*, 57, 911-917

45) Gerard, K. and Lattimer, V. (2004), "Preferences of patients for emergency services available during usual GP surgery hours: a discrete choice experiment", *Family Practice*, 22, 28-36

46) Espelid, I., Cairns, J., Askildsen, J.E., Qvist, V., Gaarden, T., Tveit, A.B. (2005), "Preferences over dental restorative materials among young patients and dental professionals", *European Journal of Oral Science*, 114, 15-21

47) Byrne, M.M., Souchek, J., Richardson, M. and Suarez-Almazor, M. (2006), "Racial/ethnic differences in preferences for total knee replacement surgery", *Journal of Clinical Epidemiology*, 59, 1078-1086

48) Ryan, M., Diack, J., Watson, V. and Smith, N. (2005), "Rapid prenatal diagnostic testing for Down syndrome only or longer wait for full karyotype: the views of pregnant women", *Prenatal Diagnosis*, 25, 1206-1211

49) Ossa, D.F., Briggs, A., McIntosh, E., Cowell, W., Littlewood, T. and Sculpher, M. (2007), "Recombinant erythropoietin for chemotherapy-related anaemia: Economic value and health-related quality-of-life assessment using direct utility elicitation and discrete choice experiment methods", *PharmacoEconomics*, 25 (3), 223-237

50) Cook, J., Whittington, D., Cahn, D.G., Johnson, F.R and Nyamete, A. (2007), "Reliability of stated preferences for cholera and typhoid vaccines with time to think in Hue, Vietnam", *Economic Inquiry*, 45 (1), 100-114

51) Gerard, K., Lattimer, V., Turnbull, J., Smith, H., George, S., Brailsford, S. and Maslin-Prothero, S. (2004), "Reviewing emergency care systems 2: measuring patient preferences using a discrete choice experiment", *Emergency Medicine Journal*, 21, 692-697

52) Osoba, D., Hsu, M., Copley-Merriman, C., Coombs, J., Johnson, F.R., Hauber, B., Manjunath, R. and Pyles, A. (2006), "Stated preferences of patients with cancer for health-related quality-of-life (HRQOL) domains during treatment", *Quality of Life Research*, 15, 273-283

53) Araña, J.E., León, C.J. and Quevedo, J.L. (2006), "The effect of medical experience on the economic evaluation of health policies. A discrete choice experiment", *Social Science and Medicine*, 63, 512-524

54) Haas, M. (2005), "The impact of non-health attributes of care on patients' choice of GP", *Australian Journal of Primary Health*, 11 (1), 40-46

55) Lloyd, A., McIntosh, E. and Price, M. (2005), "The importance of drug adverse effects compared with seizure control for people with epilepsy: A discrete choice experiment", *PharmacoEconomics*, 23 (11), 1167-1181

56) Gerard, K., Lattimer, V., Surridge, H., George, S., Turnbull, J., Burgess, A., Lathlean, J. and Smith, H. (2006), "The introduction of integrated out-of-hours arrangements in England: a discrete choice experiment of public preferences for alternative models of care", *Health Expectations*, 9, 60-69

57) Mansfield, C., Johnson, F.R. and Van Houtven, G. (2006), "The missing piece: Valuing averting behavior for children's ozone exposures", *Resource and Energy Economics*, 28, 215-228

58) Berchi, C., Dupuis, J. and Launoy, G. (2006), "The reasons of general practitioners for promoting colorectal cancer mass screening in France", *The European Journal of Health Economics*, 7, 91-98

59) Baltussen, R., Stolk, E., Chisholm, D. and Aikins, M. (2006), "Towards a multi-criteria approach for priority setting: an application to Ghana", *Health Economics*, 15, 689-696

60) Ashcroft, D.M., Seston, E. and Griffiths, C.E.M (2006), "Trade-offs between the benefits and risks of drug treatment for psoriasis: a discrete choice experiment with UK dermatologists", *British Journal of Dermatology*, 155, 1236-1241

61) Caldow, J., Bond, C., Ryan, M., Campbell, N.C., San Miguel, F., Kiger, A. and Lee, A. (2007), "Treatment of minor illness in primary care: a national survey of patient satisfaction, attitudes and preferences regarding a wider nursing role", *Health Expectations*, 10, 30-45

62) Beusterien, K.M., Dziekan, K., Flood, E., Harding, G. and Jordan, J.C. (2005), "Understanding patient preferences for HIV medications using adaptive conjoint analysis: feasibility assessment", *Value in Health*, 8 (4), 453-461

63) Hiselius, L.W. (2005), "Using choice experiments to assess people's preferences for railway transports of hazardous materials", *Risk Analysis*, 25 (5), 1199-1214

64) Dwight-Johnson, M., Lagomasino, I.T., Aisenberg, E. and Hay, J. (2004), "Using conjoint analysis to assess depression treatment preferences among low-income Latinos", *Psychiatric Services*, 55 (8), 934-936

65) Ryan, M., Netten, A., Skåtun, D. and Smith, P. (2006), "Using discrete choice experiments to estimate a preference-based measure of outcome - An application to social care for older people", *Journal of Health Economics*, 25, 927-944

66) Ryan, M., Major, K. and Skåtun, D. (2005), "Using discrete choice experiments to go beyond clinical outcomes when evaluating clinical practice", *Journal of Evaluation in Clinical Practice*, 11 (4), 328-338

67) Lancsar, E.J., Hall, J.P., King, M., Kenny, P., Louviere, J.L., Fiebig, D.G., Hossain, I., Thien, F.C.K., Reddel, H.K. and Jenkins, C.R. (2007), "Using discrete choice experiments to investigate subject preferences for preventive asthma medication", *Respirology*, 12, 127-136

68) Thompson, C.A., Foster, A., Cole, I. and Dowding, D.W. (2005), "Using social judgement theory to model nurses' use of clinical information in critical care education", *Nurse Education Today*, 25, 68-77

69) Bouma, B.J., van der Meulen, J.H.P, van den Brink, R.B.A, Smidts, A., Cheriex, E.C., Hamer, H.P., Arnold, A.E.R, Zwinderman, A.H., Lie, K.I. and Tijssen, J.G.P. (2004), "Validity of conjoint analysis to study clinical decision making in elderly patients with aortic stenosis", *Journal of Clinical Epidemiology*, 57, 815-823

70) Roux, L., Ubach, C., Donaldson, C. and Ryan, M. (2004), "Valuing the benefits of weight loss programs: An application of the discrete choice experiment", *Obesity Research*, 12 (8), 1342-1351

71) Hall, J., Fiebig, D.G., King, M.T., Hossain, I. and Louviere, J.J. (2006), "What Influences Participation in Genetic Carrier Testing? Results from a Discrete Choice Experiment", *Journal of Health Economics*, 25, 520-537

72) Mahadevia, P., Shah, S., Mannix, S., Brewster-Jordan, J., Kleinman, L., Leibman, C. and O'Dowd, L. (2006), "Willingness to pay for sensory attributes of intranasal corticosteroids among patients with allergic rhinitis", *Journal of Managed Care Pharmacy*, 12 (2),143-51

73) Walzer, S. Zweifel, P. (2007), "Willingness-to-pay for caregivers of children with asthma or wheezing conditions", *Therapeutics and Clinical Risk Management*, 3 (1), 157-166

74) Iragüen, P. and de Dios Ortúzar, J. (2004), "Willingness-to-pay for reducing fatal accident risk in urban areas: an Internet-based Web page stated preference survey", *Accident Analysis and Prevention*, 36, 513-524

75) Bishop, A.J., Marteau, T.M., Armstrong, D., Chitty, L.S., Longworth, L., Buxton, M.J. and Berlin, C. (2004), "Women and health care professionals' preferences for Down's Syndrome screening tests: a conjoint analysis study", *British Journal of Obstetrics and Gynaecology*, 111, 775-779

76) Lewis, S.M., Cullinane, F.M., Carlin, J.B. and Halliday, J.L. (2006), "Women's and health professionals' preferences for prenatal testing for Down syndrome in Australia", *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 46, 205-211

77) Christofides, N.J., Muirhead, D., Jewkes, R.K., Penn-Kekana, L. and Conco, D.N. (2005), "Women's experiences of, and preferences for, services after rape in South Africa: interview study", *British Medical Journal*, 332, 209-213

78) Wordsworth, S., Ryan, M., Skåtun, D. and Waugh, N. (2006), "Women's preferences for cervical cancer screening: a study using a discrete choice experiment", *International Journal of Technology Assessment in Health Care*, 22 (3), 344-350

79) Fraenkel, L., Constantinescu, F., Oberto-Medina, M. and Wittink, D.R. (2005), "Women's preferences for prevention of bone loss", *The Journal of Rheumatology*, 32, 1086-1092

80) Seston, E.M., Elliott, R.A., Noyes, P.R. and Payne, K. (2007), "Women's preferences for the provision of emergency hormonal contraception services", *Pharmacy World and Science*, 29, 183-189